

A FAMILY OF RAMANUJAN-ORR FORMULAS FOR $1/\pi$

JESÚS GUILLERA

ABSTRACT. We use a variant of Wan's method to prove two Ramanujan-Orr type formulas for $1/\pi$. This variant needs to know in advance the formulas for $1/\pi$ that we want to prove, but avoids the need of solving a system of equations.

1. INTRODUCTION

In [10], Wan gives a method to derive a formula for $1/\pi$ from each identity of type

$$(1) \quad G(x) = \frac{1}{\pi^2} K(a(x)) K(b(x)),$$

where G is analytic near the origin, and K is the complete elliptic integral of the first kind. The method uses only transformations of K , see [9] or [2, Chap. 19-20] for those of higher degree, and the following Legendre's identity:

$$(2) \quad -K(\sqrt{r_0})K(\sqrt{1-r_0}) + K(\sqrt{r_0})E(\sqrt{1-r_0}) + E(\sqrt{r_0})K(\sqrt{1-r_0}) = \frac{\pi}{2},$$

where E is the complete elliptic integral of the second kind [3]. The aim of this short paper is to show a variant of his method which we can use when we know in advance the formula for $1/\pi$ that we want to prove (or disprove). This variant avoids the need of solving a system of equations, because we apply the method via translation [11]. This is simpler and faster. For example, suppose that the formula

$$(3) \quad \left\{ p_0 + p_1 \left(x \frac{d}{dx} \right) + p_2 \left(x \frac{d}{dx} \right)^2 + p_3 \left(x \frac{d}{dx} \right)^3 \right\} G = \frac{1}{\pi},$$

where p_0, p_1, p_2, p_3 are known constants, holds at $x = x_0$. The translation consist in applying the same operator to the right side of (1). Then, if $a(x_0)$ and $b(x_0)$ are complementary arguments of K , the action of the operator provides a proof of the

2010 *Mathematics Subject Classification.* Primary 33E05, 33C20; Secondary 11F03, 33C75, 33F10.

Key words and phrases. Hypergeometric series; Ramanujan-Orr type formulas; Complete elliptic integral; Legendre's relation; Zeilberger's algorithm.

identity if we take into account the Legendre's identity. The following factorization:

$$(4) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{s}{2}\right)_n \left(\frac{1-s}{2}\right)_n \left(\frac{1+s}{2}\right)_n \left(1 - \frac{s}{2}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} y^n = \sum_{n=0}^{\infty} \frac{(s)_n (1-s)_n}{(1)_n^2} \left(\frac{1 - \sqrt{1-y} + \sqrt{-y}}{2}\right)^n \cdot \sum_{n=0}^{\infty} \frac{(s)_n (1-s)_n}{(1)_n^2} \left(\frac{1 - \sqrt{1-y} - \sqrt{-y}}{2}\right)^n,$$

(see [8, eq. 7.5.1.9], and [4] for a proof of it) is of type (1) for the values $s = 1/2$, $s = 1/4$, $s = 1/3$, $s = 1/6$. Another Orr-type factorization was used in [10, Sect. 4].

2. WAN'S METHOD VIA TRANSLATION

In this section we prove two couples of Ramanujan-Orr type formulas applying Wan's method via translation.

2.1. Example 1. Leting $s = 1/4$ and making $y = -4x^2(x-1)^2/(2x-1)^2$ in (4), we have

$$(5) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-4x^2(x-1)^2}{(2x-1)^2}\right)^n = \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} x^n \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} \left(\frac{x}{2x-1}\right)^n.$$

Applying the quadratic transformation

$$(6) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} x^n = \frac{1}{\sqrt{1+\sqrt{x}}} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^2}{(1)_n^2} \left(\frac{2\sqrt{x}}{1+\sqrt{x}}\right)^n, \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^2}{(1)_n^2} x^n = \frac{2}{\pi} K(\sqrt{x}),$$

to the two sums in the right side of (5), we get

$$(7) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-4x^2(x-1)^2}{(2x-1)^2}\right)^n = f(x) f\left(\frac{x}{2x-1}\right),$$

where

$$f(x) = \frac{2}{\pi} \frac{K\left(\sqrt{\frac{2\sqrt{x}}{1+\sqrt{x}}}\right)}{\sqrt{1+\sqrt{x}}}.$$

The arguments of the two elliptic integrals are complementary at

$$x_0 = \frac{1}{49} + \frac{4}{49}\sqrt{3}i, \quad y_0 = \frac{-4x_0^2(x_0-1)^2}{(2x_0-1)^2} = \frac{192}{2401}, \quad r_0 = \frac{2\sqrt{x_0}}{1+\sqrt{x_0}} = \frac{1}{2} + \frac{\sqrt{3}}{6}i.$$

This suggest that there exist formulas for $1/\pi$ with $y_0 = 192/2401$. Inspired by this, using the PSLQ algorithm, we have discovered the following ones:

$$(8) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{192}{2401}\right)^n \frac{376n^2 + 216n + 15}{2n+1} = \frac{98\sqrt{21}}{9\pi},$$

and

$$(9) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{192}{2401}\right)^n (70668n^3 - 9216n^2 + 1428n + 90) = \frac{294\sqrt{21}}{\pi}.$$

Hence, we can use Wan's method [10] to prove identity (9). However, due to the fact that we already know the formula that we want to prove (thanks to the PSLQ algorithm), we do not need constructing a system of equations. Instead, we will use the method via translation [11]; see [6] for another application of translation.

Theorem 2.1. *Formula (9) is true.*

Proof. First, by applying the operator

$$(10) \quad 90 + 1428 \left(y \frac{d}{dy} \right) - 9216 \left(y \frac{d}{dy} \right)^2 + 70668 \left(y \frac{d}{dy} \right)^3 \Big|_{y=y_0},$$

to the left side of (7), we obtain

$$(11) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{192}{2401}\right)^n (70668n^3 - 9216n^2 + 1428n + 90).$$

Then, applying the same operator, but written as

$$(12) \quad 90 + 1428 \left(\frac{y}{y'} \frac{d}{dx} \right) - 9216 \left(\frac{y}{y'} \frac{d}{dx} \right)^2 + 70668 \left(\frac{y}{y'} \frac{d}{dx} \right)^3 \Big|_{x=x_0},$$

to the right side of (7), we obtain

$$(13) \quad \frac{588\sqrt{21}}{\pi^2} \left(-K(\sqrt{r_0})K(\sqrt{1-r_0}) + K(\sqrt{r_0})E(\sqrt{1-r_0}) + E(\sqrt{r_0})K(\sqrt{1-r_0}) \right).$$

This was done automatically using the following Maple code:

```
proof:=proc()
local x0,y,w,f,g,pr1,pr,K;  K:=x->EllipticK(x);
y:=x->-4*x^2*(x-1)^2/(2*x-1)^2; w:=x->y(x)/diff(y(x),x);
f:=x->2/Pi*K(sqrt(2*sqrt(x)/(1+sqrt(x))))/sqrt(1+sqrt(x));
x0:=1/49*(1+4*sqrt(3)*I); g:=x->f(x)*f(x/(2*x-1));
pr1:=subs(x=x0,90*g(x)+1428*w(x)*diff(g(x),x)
-9216*w(x)*diff(w(x)*diff(g(x),x),x)
+70668*w(x)*diff(w(x)*diff(w(x)*diff(g(x),x),x),x));
pr:=combine(simplify(combine(simplify(expand(
simplify(evalc(simplify(pr1))))),radical)),radical);
return pr;
end;
```

Copy and paste this code in a Maple session. Executing the procedure `proof()`; gives the output (13). Finally, taking into account the Legendre's identity (2), formula (9) follows. $\square$

The following theorem implies that the formulas (8) and (9) are equivalent.

Theorem 2.2. *The identity*

$$(14) \quad \sum_{n=0}^{\infty} B(n, s) z^n \left[\frac{s(s^2 - 1)(s - 2) + 4(1 + 2s - 2s^2)n + 8(1 + s - s^2)n^2}{2n + 1} - \frac{8(1 - z)}{z} n^3 + 12n^2 \right] = 0,$$

where

$$B(n, s) = \frac{\left(\frac{s}{2}\right)_n \left(\frac{1-s}{2}\right)_n \left(\frac{1+s}{2}\right)_n \left(1 - \frac{s}{2}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3},$$

holds.

Proof. We can prove it automatically by computer [7] writing in a Maple session

```
with(SumTools[Hypergeometric]):
print(op=Zeilberger(v(n,s),s,n,S)[1]):
w:=(n,s)->subs({p=n,q=s},Zeilberger(v(p,q),q,p,S)[2]):
print(w0=w(0,s));
```

where $v(n, s)$ is the function inside the sum of (14), and S is the operator such that $S v(n, s) = v(n, s+1)$. The output $op=1$ (independent of S), means that

$$1 \cdot v(n, s) = v(n, s) = w(n+1, s) - w(n, s).$$

Then, as $w(0, s) = 0$, identity (14) is due to telescoping cancelation when we sum for $n \geq 0$. $\square$

Remark 2.3. It is possible to check the proof of (8) and (9) in a completely automatic way because a computer can verify that:

- (1) Both sides of (7) satisfy the same differential equation.
- (2) The action of the operator (10) to the left side of (7) gives (11). This is trivial, even for human.
- (3) The action of the operator (12) to the right side of (7) gives (13). See the procedure `proof()`.
- (4) Identity (14) is true (see the Maple code above).

Observe that these steps are elementary in the sense that they do not require the use of modular functions. Observe in addition that the value of τ , such that

$$\frac{K(\sqrt{1-r_0})}{K(\sqrt{r_0})} = -2i\tau, \quad q = e^{2\pi i\tau},$$

corresponds to the value of r_0 of the above proof, is not a quadratic irrational (in fact it seems to be transcendent).

2.2. **Example 2.** Letting $s = 1/4$ and replacing x with $x/(x-1)$ in (5), we have

$$(15) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-4x^2}{(x^2-1)^2}\right)^n = \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} \left(\frac{x}{x-1}\right)^n \cdot \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} \left(\frac{x}{x+1}\right)^n,$$

Then, using the known transformation

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{(1)_n^2} \left(\frac{x}{x-1}\right)^n = \sqrt[4]{1-x} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^2}{(1)_n^2} \left(\frac{1}{2} - \frac{1}{2}\sqrt{1-x}\right)^n,$$

and the quadratic transformations (6) and

$$K(x) = \frac{1}{1+x} K\left(\frac{2\sqrt{x}}{1+x}\right),$$

we deduce that

$$(16) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-4x^2}{(x^2-1)^2}\right)^n = \frac{4}{\pi^2} \frac{\sqrt[4]{1-x}}{(1+g(x))\sqrt{1+h(x)}} K\left(\frac{2\sqrt{g(x)}}{1+g(x)}\right) K\left(\sqrt{\frac{2h(x)}{1+h(x)}}\right),$$

where

$$g(x) = \sqrt{\frac{1}{2} - \frac{1}{2}\sqrt{1-x}}, \quad h(x) = \sqrt{\frac{x}{x+1}}.$$

The arguments of the elliptic K integrals in (16) are complementary at

$$x_0 = \frac{85\sqrt{41} - 529}{128}, \quad y_0 = -\frac{2^{14}}{23^4} = -\left(\frac{128}{529}\right)^2, \quad r_0 = \frac{2h(x_0)}{1+h(x_0)} = \frac{33 - 5\sqrt{41}}{2},$$

where $y(x) = -4x^2/(x^2-1)^2$. This suggest that there exist formulas for $1/\pi$ with $y_0 = -2^{14} \cdot 23^{-4}$. With the help of the PSLQ algorithm we have found

$$(17) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-2^{14}}{23^4}\right)^n \frac{6970n^2 + 4037n + 280}{529(2n+1)} = \frac{\sqrt{23}}{3\pi},$$

and the contiguous formula

$$(18) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-2^{14}}{23^4}\right)^n \frac{-296225n^3 - 24576n^2 + 53002n + 4200}{3174} = \frac{\sqrt{23}}{\pi}.$$

Theorem 2.4. *Formulas (17) and (18) are true.*

Proof. First, we apply the operator

$$(19) \quad 4200 + 53002 \left(y \frac{d}{dy} \right) - 24576 \left(y \frac{d}{dy} \right)^2 - 296225 \left(y \frac{d}{dy} \right)^3 \Big|_{y=y_0},$$

to the left side of (16). Then, we use the Maple code

```
proof:=proc()
local x0,y,w,g,h,f,pr1,pr,prr,K; K:=x->EllipticK(x);
y:=x->-4*x^2/(x^2-1)^2; w:=x->y(x)/diff(y(x),x);
x0:=-529/128+85/128*sqrt(41);
g:=x->sqrt(1/2-1/2*sqrt(1-x)); h:=x->sqrt(x/(x+1));
f:=x->4/Pi^2*(1-x)^(1/4)*1/((1+g(x))*sqrt(1+h(x)))
*K(2*sqrt(g(x))/(1+g(x)))*K(sqrt(2*h(x)/(1+h(x))));
pr1:=subs(x=x0,4200*f(x)+53002*w(x)*diff(f(x),x)
-24576*w(x)*diff(w(x)*diff(f(x),x),x)
-296225*w(x)*diff(w(x)*diff(w(x)*diff(f(x),x),x),x));
pr:=simplify(combine(simplify(expand(simplify(evalc(
simplify(pr1))))),radical));
prr:=simplify(radnormal(combine(radnormal(pr),radical)));
return prr;
end;
```

to apply the same operator, but written as

$$(20) \quad 4200 + 53002 \left(\frac{y}{y'} \frac{d}{dx} \right) - 24576 \left(\frac{y}{y'} \frac{d}{dx} \right)^2 - 296225 \left(\frac{y}{y'} \frac{d}{dx} \right)^3 \Big|_{x=x_0},$$

to the right side of (16). Executing the procedure and using the Legendre's identity, we prove (18), and from it we deduce (17) by using the identity (14). $\square$

3. RAMANUJAN-ORR TYPE FORMULAS AND THE PSLQ ALGORITHM

By the PSLQ algorithm we have discovered the formulas

$$(21) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \frac{1}{7^{4n}} \frac{1920n^2 + 1072n + 55}{2n + 1} = \frac{196\sqrt{7}}{3\pi},$$

and

$$(22) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n \left(\frac{1}{6}\right)_n \left(\frac{5}{6}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{3}{5}\right)^{6n} \frac{133n^2 + 79n + 6}{2n + 1} = \frac{625}{32\pi}.$$

Hence, we believe that it should be possible to prove (21) and (22) from (4), with $s = 1/4$ and $s = 1/3$ respectively. In addition we find curious that these formulas look similar to the following challenging series for $1/\pi^2$:

$$(23) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{(1)_n^5} \frac{1}{7^{4n}} (1920n^2 + 304n + 15) = \frac{56\sqrt{7}}{\pi^2},$$

and

$$(24) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n \left(\frac{1}{6}\right)_n \left(\frac{5}{6}\right)_n}{(1)_n^5} \left(\frac{3}{5}\right)^{6n} (532n^2 + 126n + 9) = \frac{375}{\pi^2}.$$

Formula (23) was conjectured in [5], and formula (24) is joint with G. Almkvist, and was conjectured in [1]. The following example of formula for $1/\pi$:

$$(25) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{10}\right)_n \left(\frac{3}{10}\right)_n \left(\frac{7}{10}\right)_n \left(\frac{9}{10}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \frac{1}{2^{6n}} \frac{2100n^2 + 1160n + 63}{2n + 1} = \frac{200}{\pi},$$

is such that there is no known Orr-type factorization from which we can derive it.

As an example of application of the PSLQ algorithm, we explain how we have discovered the formula (17): Let

$$t(j) = \sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-2^{14}}{23^4}\right)^n \frac{n^j}{2n + 1}.$$

Then, looking for integer relations among the quantities

$$\frac{1}{\pi^2}, \quad t(0)^2, \quad t(1)^2, \quad t(2)^2, \quad t(0)t(1), \quad t(0)t(2), \quad t(1)t(2),$$

we guess that

$$\begin{aligned} \frac{6436343}{\pi^2} = & 705600 t(0)^2 + 146676321 t(1)^2 + 437228100 t(2)^2 \\ & + 20346480 t(0)t(1) + 35128800 t(0)t(2) + 506482020 t(1)t(2), \end{aligned}$$

and taking the square root, we have

$$\frac{529\sqrt{23}}{\pi} = 3(6970 t(2) + 4037 t(1) + 280 t(0)).$$

This is interesting because as we have explained before, when we guess a formula corresponding to complementary arguments of the elliptic integrals of an Orr-type factorization, then it is simple to prove.

ACKNOWLEDGEMENT

I thank Wadim Zudilin for encouraging me to extend these investigations with the use of modular equations in alternative bases.

REFERENCES

- [1] G. ALMKVIST, AND J. GUILLERA, Ramanujan-like series for $1/\pi^2$ and String theory. *Exp. Math.* **21**, (2012), 223-234.
- [2] B.C. BERNDT, *Ramanujan's Notebooks, Part III* (Springer-Verlag, New York, 1991).
- [3] J.M. BORWEIN AND P.B. BORWEIN, Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity, *Canad. Math. Soc. Series Monographs Advanced Texts* (Jonh Wiley, New York, 1987).
- [4] H.H. CHAN AND Y. TANIGAWA, A generalization of a Brafman-Bailey type identity. *Proceedings of the Amer. Math. Soc.* **143**, (2015), 185-195.

- [5] J. GUILLERA, About a new kind of Ramanujan type series, *Experimental Mathematics* **12**, (2003), 507-510.
- [6] J. GUILLERA AND W. ZUDILIN, Ramanujan-type formulae for $1/\pi$. The art of translation, in *The Legacy of Srinivasa Ramanujan*, B.C. Berndt & D. Prasad (eds.), *Ramanujan Math. Soc. Lecture Notes Series* **20** (2013), 181–195
- [7] M. PETKOVŠEK, H.S. WILF, AND D. ZEILBERGER, A=B. Wellesley, Massachusetts, A K. Peters, Ltd., 1996.
- [8] A.P. PRUDNIKOV, YU.A. BRYCHKOV, O.I. MARICHEV, Integrals and series Vol.3. More Special Functions. *New York-London: Gordon & Breach*, 1990.
- [9] R. VIDŪNAS, Algebraic transformations of Gauss hypergeometric functions. *Funkcial. Ekvac.* **52** (2009), 139–180.
- [10] J. WAN, Series for $1/\pi$ Using Legendre’s Relation, *Integral Transforms and Special Functions* **25**, p. 1-14, (2014).
- [11] W. ZUDILIN, Lost in translation, in *Advances in Combinatorics*, Waterloo Workshop in Computer Algebra W80 (May 26–29, 2011), I. Kotsireas and E.V. Zima (eds.) (2013), Springer, 287–293.

4. ADDENDUM

From (15), we can arrive at

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-4x^2}{(x^2-1)^2}\right)^n = \frac{4}{\pi^2} \frac{\sqrt[4]{1-x}}{\sqrt{1+h(x)}} K\left(\sqrt{g(x)}\right) K\left(\sqrt{\frac{2h(x)}{1+h(x)}}\right),$$

(26) where $g(x) = \frac{1}{2} - \frac{1}{2}\sqrt{1-x}$, and $h(x) = \sqrt{\frac{x}{x+1}}$.

From (26), and in the same way used to derive the previous formulas in this paper, we can prove the following “divergent” series for $1/\pi$:

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-2^{14}}{7^4}\right)^n (-600 - 7518n - 24576n^2 - 18785n^3) \stackrel{?}{=} \frac{98\sqrt{7}}{\pi},$$

(27) $\sum_{n=0}^{\infty} \frac{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n^3} \left(\frac{-2^{14}}{7^4}\right)^n \frac{120 + 1273n + 2210n^2}{2n+1} \stackrel{?}{=} \frac{49\sqrt{7}}{\pi}.$

Since these last two formulas are “divergent” their “upside-down” are convergent, and inspired by [1], we have guessed that

$$\sum_{n=1}^{\infty} \frac{\left(\frac{1}{2}\right)_n (1)_n^3}{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n} \left(\frac{-7^4}{2^{14}}\right)^n \frac{-600 + 7518n - 24576n^2 + 18785n^3}{7^4 n^3} \stackrel{?}{=} 2L_{-7}(2) - 1,$$

(28) $\sum_{n=1}^{\infty} \frac{\left(\frac{1}{2}\right)_n (1)_n^3}{\left(\frac{1}{8}\right)_n \left(\frac{3}{8}\right)_n \left(\frac{5}{8}\right)_n \left(\frac{7}{8}\right)_n} \left(\frac{-7^4}{2^{14}}\right)^n \frac{120 - 1273n + 2210n^2}{7^4 (-2n+1)n^3} \stackrel{?}{=} L_{-7}(2).$

A geometric interpretation (as a volume) of the arithmetical constant $L_{-7}(2)$ is given in [2].

REFERENCES

- [1] J. GUILLERA AND M. ROGERS, Ramanujan series upside-down. *Journal of the Australian Math. Society* **97**, (2014), 78-106.
- [2] D.H. BAILEY, J.M. BORWEIN, D. BROADHURST & W. ZUDILIN, Experimental mathematics and mathematical physics, in *Gems in Experimental Mathematics*, T. Amdeberhan, L.A. Medina, and V.H. Moll (eds.), *Contemp. Math.* **517**, (2010), Amer. Math. Soc., 41–58

AV. CESÁREO ALIERTA, 31 ESC. IZ. 4^o-A, ZARAGOZA (SPAIN)

E-mail address: jguillera@gmail.com